

The role of preovulatory estradiol in fertility

George A. Perry and Emmalee J. Northrop

South Dakota State University, Department of Animal Science

INTRODUCTION

The most efficient and cost-effective method for genetic improvement of economically important traits in the beef industry is artificial insemination (AI). Estrous synchronization and AI remain the most important and widely applicable reproductive biotechnology available for cattle (Seidel, 1995). However, the time and labor required to detect spontaneous estrus in beef cattle has been a deterrent to the widespread utilization of AI (Britt, 1987). Therefore, the development of procedures that facilitate high pregnancy rates at fixed-time insemination could expedite genetic progress.

At the initiation of estrus, gonadotropin-releasing hormone (GnRH) is secreted from the hypothalamus at a sufficient amplitude and frequency to stimulate a surge of luteinizing hormone (LH) to be released from the anterior pituitary to cause ovulation (Karsch et al., 1997). However, even when estrus does not occur a GnRH agonist is capable of causing ovulation of dominant bovine follicles (Ryan et al., 1998), and the luteal tissue that forms is capable of undergoing prostaglandin (PG)-induced luteal regression on days 6 or 7 (GnRH-PG protocol, Twagiramungu et al., 1995). The addition of a second injection of GnRH 48 hours after PG has been utilized to induce ovulation and eliminate the need for estrus detection. The second GnRH injection in this method resulted in induced ovulation of a wide range of follicle sizes (Perry et al., 2005).

A recent study involving the reciprocal embryo transfer of embryos to and from cows induced to ovulate either a large or small follicle with GnRH revealed some interesting results about the factors affecting fertility (Atkins et al., 2010, submitted). While ovulatory follicle size and serum concentrations of estradiol were highly correlated ($r = 0.49$; $P < 0.0001$), both concentrations of estradiol and follicle size had independent positive effects on fertilization success. Furthermore, a positive relationship between estrus expression at time of insemination and pregnancy success has been established in both beef and dairy cattle (Vasconcelos et al., 2001; Perry et al., 2005; Lopes et al., 2007; Perry et al., 2007). Among beef cows and heifers, analysis of more than 10,000 animals from 22 different studies, utilizing one of the top five recommended fixed-time AI protocols, indicated that $57.1\% \pm 3.5\%$ of animals exhibited standing estrus prior to fixed-time AI, and those animals that exhibited estrus averaged 27% greater pregnancy success compared to animals that did not (Richardson et al., 2016). Among high producing dairy cows, current rates of estrus detection are not sufficient for efficient reproductive management (Nebel and McGilliard, 1993; Senger, 1994; Dransfield et al., 1998; Lopez et al., 2004; Peralta et al., 2005).

In addition to the use of supplemental GnRH, several studies have investigated the role of supplemental estrogen on fertility (Cerri et al., 2004; Souza et al., 2005; Brusveen et al., 2009). Estradiol has a pivotal role on the initiation of estrus and the ovulatory cascade (Allrich, 1994). Preovulatory estradiol concentrations also play a critical role in sperm transport, regulation of the uterine environment, and embryo survival. Recent data from our laboratory has reported that some follicles that are ovulated in response to a GnRH injection are incapable of producing sufficient estradiol to initiate standing estrus while others are not.

ESTRADIOL PRODUCTION

A capillary network present in the thecal layer is responsible for supplying cholesterol for steroid production. Cholesterol is delivered to the theca interna cells where it is converted into androstenedione. Androstenedione is then transported to the granulosa cells where it is converted into estradiol by the aromatase enzyme (two cell – two gonadotropin concept, Fortune and Quirk, 1988). Thus, increased estradiol production by preovulatory follicles depends on the enhanced ability of the theca interna cells to produce androstenedione and the enhanced ability of granulosa cells to convert androstenedione to estradiol (Ginther et al., 1996). Estradiol production can also be increased by production of pregnenolone by granulosa cells, which can be converted to androstenedione by the theca interna cells, which in turn can be converted to estradiol by the granulosa cells (Fortune, 1986).

Estradiol production is further increased due to an increase in luteinizing hormone (LH) pulse frequency (Fortune et al., 2001). Pituitary gonadotropes produce and secrete LH, and the response of gonadotropes to GnRH is directly correlated with the number of GnRH receptors on the cell surface (Wise et al., 1984). Expression of the GnRH receptor can be regulated by estradiol, progesterone, and GnRH itself (Gregg et al., 1991; Wu et al., 1994; Looper et al., 2003). Increased sensitivity of gonadotropes to GnRH and increased expression of GnRH receptors occurs prior to the increase in concentrations of estradiol (Turzillo et al., 1994). Therefore, the decrease in progesterone before proestrus may be important in initiating the increase in LH secretion that leads to the increase in estradiol production. As progesterone decreases to basal concentrations at luteolysis, GnRH pulse frequency increased (Chabbert-Buffeta et al., 2000) and that increase in GnRH stimulated greater expression of GnRH receptors (Nett et al., 2002). Therefore, the ability of some dominant follicles to produce sufficient concentrations of estradiol to induce estrus may be a result of increased pituitary sensitivity to GnRH following a decrease in progesterone.

THE ROLE OF ESTRADIOL IN REGULATING FERTILITY

Estrus

Initiation of estrus occurs due to increased circulating concentrations of estradiol at a time when progesterone is not present (Allrich, 1994). Estrus refers to the period of time when a female is sexually receptive. Standing estrus refers to the visual sign that a female will stand to be

mounted by a male or another female. Among cattle, concentrations of estradiol peak approximately 36 hours before ovulation (Chenault et al., 1975), and this increased preovulatory concentrations of estradiol have been correlated with increased pregnancy success (Perry et al., 2005). The ability of estradiol to synchronize estrus and ovulation and its impact on overall fertility has been the focus of many investigations.

Pre-ovulatory follicular environment

The endocrine microenvironment of a preovulatory follicle has been correlated with oocyte quality and its ability to undergo germinal vesicle breakdown (McNatty et al., 1979). This microenvironment is unique relative to surrounding nonovulatory follicles and is important in the preparation of follicular cells for luteinization and secretion of progesterone (McNatty et al., 1975). Follicles containing oocytes that were more capable of being fertilized and developing to the blastocyst stage embryos contained less progesterone (Hazeleger et al., 1995), three to eight fold greater aromatase activity, greater amounts of the α subunit of inhibin (Driancourt et al., 1998), and greater concentrations of estradiol (Mermillod et al., 1999). In addition, the ability of human oocytes to develop into embryos increased when they were collected from follicles having greater follicular fluid concentrations of estradiol compared to oocytes collected from follicles that had lower concentrations of estradiol (Teissier et al., 2000). Follicular secretion of estradiol may also affect the establishment and maintenance of pregnancy through altering sperm transport and the uterine environment.

Sperm Transport

Sperm transport through the female reproductive tract is affected by estradiol and is optimized at estrus or when females are under the influence of estrogen (Hawk, 1983). Ovariectomized ewes (Allison and Robinson, 1972) and rabbits (Noyes et al., 1959) required estradiol for appropriate sperm transport through the reproductive tract. The dependence of sperm transport on the effects of estradiol is likely associated with changes in uterine pH as considerable evidence exists that pH regulates sperm motility and preserves viability (Wong et al., 1981). In cattle, there is a transient decline in uterine pH from 7.0 to 6.7 around the initiation of standing estrus (Elrod and Butler, 1993; Perry and Perry, 2008a). The importance of this decrease in uterine pH at the initiation of standing estrus to increase fertility has been reported in lactating beef cows (Bolzenius et al., 2016). Cows that initiated standing estrus had decreased uterine pH, 6.78, and increased pregnancy success, 52%, compared to cows that did not initiate standing estrus but were induced to ovulate, 6.96 and 38%, respectively.

Importantly, work from our laboratory indicates that cattle which exhibited standing estrus and possessed elevated concentrations of estradiol prior to insemination had increased pregnancy success compared to cattle that did not exhibit standing estrus (Perry et al., 2005; Perry et al., 2007). This increased pregnancy success among cows that exhibited standing estrus is likely due to increased sperm transport to the site of fertilization (Larimore et al., 2015). Subsequent work

by our laboratory (Perry and Perry, 2008a, b) has focused on the effects of estradiol on uterine pH of cattle where supplementation of estradiol decreased uterine pH to a level similar to cows in standing estrus. Similar evidence can be found in the mouse, where Wang and others (Wang et al., 2003) reported Na⁺/H⁺ exchanger (NHE) isoforms 1, 2 and 4 were present in mouse endometrial epithelial during the luteal phase and regulated uterine pH by extruding H⁺ into the uterine lumen to increase endometrial secretion of HCO₃⁻ and uterine pH. Among cows that exhibited estrus, expression of mRNA for NHE1 was decreased at 48 hours compared to hour 0, mRNA for NHE2 was decreased at 12, 24, 36, 48 and 60 hours compared to hour 0, and mRNA for NHE4 was decreased at 36, 48 and 60 hours compared to hour 0. Among cows that did not exhibit estrus, expression of NHE1, NHE2 and NHE4 did not change over time (Bolzenius et al., 2016).

Following estrus, cattle exhibit a rapid increase in uterine pH from 6.7 (day 0) to 7.0 prior to the time of ovulation (Perry and Perry, 2008a, b), and the neutral pH of 7 is maintained through the luteal phase of the cycle (Elrod et al., 1993). As extracellular pH increased up to 7.5, sperm motility increased in demembrated bull (Goltz et al., 1988), membrane intact bull (Contri et al., 2013), and human (Giroux-Widemann et al., 1991) sperm. Increased sperm motility appeared necessary for sperm to penetrate the viscous oviductal mucus and the cumulus matrix (Suarez and Dai, 1992) as well as the zona pellucida for fertilization (Stauss et al., 1995). Furthermore, capacitation (a process necessary for sperm egg binding) is initiated through an increase in intracellular pH (Fraser et al., 1993). Following capacitation and initiation of hyperactive motility, sperm must undergo the acrosome reaction in order to fertilize an ovum (Yanagimachi and Bhattacharyya, 1988). Sperm did not undergo the acrosome reaction when an increase in intracellular pH was prevented (Parrish et al., 1989; Zeng et al., 1996). Estradiol induced a decrease in uterine pH at the initiation of standing estrus, which may initially decrease sperm motility, inhibit capacitation, and the acrosome reaction thereby increasing sperm longevity. The rapid rise in uterine pH prior to ovulation is necessary for increased sperm motility, and to initiate capacitation and the acrosome reaction in order to be able to fertilize the oocyte.

Fertilization

A recent review by Santos (2004) reported fertilization failure in lactating beef and dairy cows was as high as 45%. A recent study in beef cows, in which embryos were flushed from cows that had high or low concentrations of estradiol at time of fixed-time AI revealed that cows with greater concentrations of estradiol at the GnRH-induced ovulation were more likely to yield a fertilized embryo than an unfertilized oocyte (Jinks et al., 2013). Data from our laboratory has reported that animals that exhibited estrus prior to fixed-time AI had increased accessory sperm numbers and improved embryo quality compared to animals that did not exhibit estrus (Larimore et al., 2015). Although accessory sperm are not involved in fertilization, they represent sperm that were able to access the oviduct, undergo capacitation and the acrosome reaction, recognize and bind to the oocyte, and partially penetrate the zona pellucida (Dalton et al., 2006). The number of

accessory sperm trapped in the zona pellucida has been positively associated with fertility (Hunter and Wilmut, 1984; Hawk and Tanabe, 1986; DeJarnette et al., 1992; Nadir et al., 1993), and are thought to be an indirect measure of both sperm transport and the availability of healthy competent sperm competing for fertilization (DeJarnette et al., 1992). These studies indicate the important role preovulatory estradiol concentrations have on sperm transport and fertilization success.

Oviduct and uterine gene expression changes

Ovarian estradiol has been reported to play an important role in establishing the timing of uterine receptivity (Ozturk and Demir, 2010). More specifically, estradiol likely plays a direct role in regulating oviductal secreted glycoproteins (Buhi, 2002) and in the regulation of the biological clock in the uterus (Nakamura et al., 2005). Whether or not an animal is exposed to elevated concentrations of estradiol impacts gene expression changes within the endometrium (Bridges et al., 2012). Specifically, estrus expression at time of AI has been shown to influence endometrium and corpus luteum gene expression. Cows that showed estrus have been reported to have increased expression of genes related to the maternal immune system and adhesion molecules, and downregulation of genes associated with prostaglandin synthesis and CL maintenance (Davoodi et al., 2016).

The TGF- β superfamily is involved in changes in the endometrium, development of the placenta, and maintenance of the pregnancy (Jones et al., 2006). At estrus there is also up-regulation of several genes involved in the remodeling of the extracellular matrix (Bauersachs et al., 2005), and changes in expression of inhibin α subunit (a member of the TGF- β signaling pathway) in the bovine intercaruncular area suggests it is involved in coordinating endometrial remodeling (Ishiwata, 2003 #1582). Furthermore, a recent study in cattle examined the role of high and low preovulatory estradiol concentrations on regulation of progesterone and estradiol receptors later in the estrous cycle. The authors reported that progesterone receptors in the deep glandular epithelium as well as ER α receptor mRNA in the endometrium were up-regulated in cows with elevated preovulatory estradiol concentrations (Bridges et al., 2012).

Milk protein (SERPINA14) is expressed in the endometrium of ruminants during pregnancy and is likely involved in nutrition of the embryo/fetus, growth of the embryo/fetus, and suppression of the maternal immune system (Ing and Roberts, 1989). Expression of SERPINA14, as determined by Real-time RT-PCR, was greatest on the day of estrus and expression was also up-regulated after stimulation with estradiol. Furthermore, detection of SERPINA14 protein revealed increased protein concentrations on the day of estrus and was localized to the glandular epithelium (Ulbrich et al., 2009). This suggests a possible preparatory role for preovulatory estradiol in establishing a uterine environment conducive to pregnancy. Estradiol induced progesterone (Zelinski et al., 1980) and oxytocin (Lamming and Mann, 1995) endometrial receptors in ruminants. Expression of uterine oxytocin and steroid receptors (estradiol receptor α and nuclear progesterone receptor) changed throughout the estrous cycle (Robinson et al.,

2001), and they play a vital role in regulating the uterine environment. Therefore, estradiol may not only have a direct effect on regulating the uterine environment, but may also have an indirect role by regulating response to progesterone through regulation of the progesterone receptor.

Subsequent Progesterone effects

The endocrine microenvironment of a preovulatory follicle is also unique relative to surrounding nonovulatory follicles and is important for preparation of follicular cells for luteinization and secretion of progesterone (McNatty et al., 1975). McNatty et al. (1979) suggested that development of a normal CL depends upon a follicle meeting the following criteria: 1) an adequate number of granulosa cells, 2) an adequate number of LH receptors on the granulosa and thecal cells, and 3) granulosa cells capable of synthesizing adequate amounts of progesterone following luteinization. Within granulosa cells, estradiol is reported to cause: 1) increased cellular proliferation (Goldenberg et al., 1972; Parrott and Skinner, 1998; Dupont et al., 2000), 2) formation of gap junctions (Merk et al., 1972; Burghardt and Anderson, 1981), 3) increased stimulatory action of FSH on aromatase activity (Adashi and Hsueh, 1982; Zhuang et al., 1982; Reilly et al., 1996), 4) enhanced stimulation of progesterone synthesis following gonadotropin stimulation (Welsh et al., 1983; Fanjul et al., 1984), and 5) enhanced acquisition of LH receptors (Kessel et al., 1985; Farookhi and Desjardins, 1986; Wang and Greenwald, 1993). Furthermore, luteinized granulosa cells secreted increased amounts of progesterone when they were collected from follicles having increased concentrations of estradiol compared to granulosa from follicles that had lower concentrations of estradiol (McNatty et al., 1979).

The relationship between pregnancy success and circulating concentrations of progesterone during early pregnancy in cattle is equivocal, as luteal secretion of progesterone is required for the survival of the embryo/fetus (McDonald et al., 1952). Several studies have reported elevated concentrations of progesterone in pregnant cows compared to nonpregnant cows beginning as early as day 4 (Butler et al., 1996) or day 6 (Henricks et al., 1971; Erb et al., 1976) after insemination. Furthermore, cows that had an earlier rise in progesterone had embryos that were more advanced developmentally, produced more interferon τ (INF- τ) and were capable of inhibiting the PGF_{2 α} release on day 16 after breeding (Kerbler et al., 1997; Mann et al., 1998; Mann et al., 1999; Mann and Lamming, 2001). Similarly, cows supplemented with progesterone during early gestation had advanced endometrial expression of several genes associated with uterine secretions and conceptus development (Forde et al., 2009; Forde et al., 2011). It is suspected that progesterone induces changes in endometrial gene expression that leads to changes in uterine histotroph composition (Spencer et al., 2008). Bartol et al. (1981) determined that protein accumulation within the uterine lumen is related to length of progesterone stimulation. However, direct supplementation of progesterone following insemination has produced mixed results. Some studies have reported a 10 to 60% increase in pregnancy rates following progesterone supplementation (Johnson et al., 1958; Robinson et al., 1989; Macmillan and Peterson, 1993), but in other studies using sheep (Diskin and Niswender, 1989; Nephew et

al., 1994) and cattle (Walton et al., 1990; Van Cleef et al., 1991) progesterone supplementation had no benefit on pregnancy success. A consistent characteristic among animals in which progesterone supplementation increased pregnancy rate was that the animals appeared to have fertility problems, since control animals had unusually low pregnancy rates.

Work by Atkins et al. (2013) reported that estradiol concentrations at GnRH-induced ovulation affected day 27 pregnancy rate of recipient cows independent of progesterone on day 7. Furthermore, work from our laboratory has reported when ovulatory follicle size was controlled there was no difference in day 10 CL weight, circulating concentrations of progesterone, or expression of luteal steroidogenic enzymes between cows that exhibited standing estrus and cows that did not (Fields et al., 2012). Therefore, improved pregnancy success among cattle with elevated preovulatory concentrations of estradiol are likely independent of the impact of progesterone on the uterine environment during the subsequent estrous cycle.

Uterine environment.

The uterine environment plays a critical role in early embryo development, recognition of pregnancy, elongation, and attachment. During the estrous cycle changes occur in the endometrium regarding its composition and differentiation status. These changes are mainly regulated by estradiol, progesterone, and oxytocin (Spencer and Bazer, 2004). The timing of changes in the uterine environment is of critical importance to embryo survival. In cattle, synchrony between the embryo and the uterus must be ± 24 hours (Hasler, 2001). Ovarian estradiol plays an important role in establishing the timing of uterine receptivity (Ozturk and Demir, 2010).

CLOCK genes regulate biological oscillations and the transcription of other genes in a tissue specific manner in response to changes in nutrient status, day length, and possibly other environmental influences (Albrecht, 2006). Decreased litter size, lower frequency of successful matings, and increased cycle length have been reported in *Clock* ^{Δ 19} (mice in which CLOCK is not functional) female mice (Chappell et al., 2003). When *Clock* ^{Δ 19} female mice were treated hormonally to induce ovulation and mated with fertile males, *Clock* ^{Δ 19} females produced fewer embryos than control females (Kennaway, 2005). It is possible from these results that defects in circadian clock gene expression at the level of the uterus may disrupt the synchrony between the uterine environment and the developing embryo. He and co-workers (2007) reported that Period 1 mRNA (part of the positive feedback loop of CLOCK) was expressed in the uterus of diestrus rats and that both estradiol and progesterone stimulated uterine Period 1 mRNA expression in the uterus of ovariectomized rats. Furthermore, Nakamura and co-workers (2005) reported that estradiol altered the normal circadian rhythms in the uterus. Therefore, the circadian clock genes may influence fertility at the level of the uterus.

The uterine histotroph is composed of nutrients, growth factors, proteins, glucose, immunosuppressive agents, enzymes, and ions. It is secreted by the endometrium, and is necessary for early conceptus growth/survival (Geisert et al., 1992; Gray et al., 2001). Any changes can greatly influence early embryonic viability. In particular, glucose is a major fuel source used by the conceptus for growth and development. Work from our laboratory found that animals that showed estrus during a fixed-time AI program had greater glucose in the uterine luminal fluid compared to no estrus animals (unpublished). A subsequent study determined that animals that exhibit estrus had increased mRNA expression of specific glucose transporters in the endometrium. In both caruncular and intercaruncular endometrial tissue, both SLC2A1 and SLC5A1 mRNA abundance was increased in animals that exhibited estrus ($P \leq 0.05$). This may serve as a possible mechanism to deliver glucose into the uterus where it can be utilized by the developing conceptus.

When an injection of estrogen, corresponding with initiation of estrus, was omitted in ovariectomized hormone-supplemented ewes, embryo survival following embryo transfer (Miller and Moore, 1976), uterine weight, uterine protein, RNA to DNA ratio, and the rate of protein synthesis were decreased (Miller et al., 1977). Elevated concentrations of estradiol did not impact protein to DNA, RNA to DNA ratios or in-vitro rate of protein synthesis in intercaruncular and caruncular endometrium on the day of estrus, but increased all measures from day 2 to 4 after estrus (Miller and Moore, 1983). Furthermore, cows that exhibit standing estrus around the time of fixed-time insemination had increased preovulatory concentrations of estradiol (Perry et al., 2005; Perry and Perry, 2008a, b), decreased uterine pH (Perry and Perry, 2008a, b), and increased pregnancy success (Perry et al., 2005) compared to cows that did not exhibit standing estrus. Consequently, preovulatory concentrations of estradiol may play a major role in the uterine environment and the establishment and maintenance of pregnancy.

Early embryo development and survival

Production of a glycoprotein, known as oviductal glycoprotein (OGP) also occurs during estrus by the non-ciliated epithelial cells of the oviduct from the late follicular phase through early embryo development (Buhi, 2002). When cyclic ewes were ovariectomized production of OGP was abolished, but administration of estradiol benzoate restored OGP production (Sutton et al., 1986). Oviductal glycoprotein has been reported to be present on the head, midpiece (King and Killian, 1994), and tail (Abe et al., 1995) of bovine sperm in oviductal fluid, and OGP may be instrumental in fertilization success and early embryo development. *In vitro* matured oocytes cultured in media containing varying concentrations of OGP resulted in increased cleavage rates and blastocyst formation following IVF in a dose-dependent manner (Hill et al., 1996).

When an injection of estrogen, corresponding with initiation of estrus, was omitted in ovariectomized hormone-supplemented ewes, embryo survival following embryo transfer was decreased (Miller and Moore, 1976). In a recent study from our laboratory ovariectomized cows

received exogenous hormones to mimic the luteal phase and luteolysis and then received either estradiol cypionate (ECP), estradiol benzoate (EB) or no treatment (CON) according to Latin Square assignment to mimic a preovulatory period. Ovulation was simulated with an injection of gonadotropin releasing hormone (GnRH, 100 µg; day 0), and embryos were transferred on day 7. Pregnancy was maintained by exogenous progesterone supplementation. Expression of interferon stimulated genes (ISG) was used as a marker for pregnancy on day 17, 19, 21, and 28 and pregnancy-specific protein B (PSPB) was used as a marker for pregnancy on day 22-28. Transrectal ultrasonography was used on day 29 to diagnose pregnancy. Cows that received preovulatory estradiol exposure had greater embryonic survival and pregnancy establishment compared to control animals (4%, 29%, and 21% for CON, EB, and ECP, respectively; (Madsen et al., 2015). Additionally, preovulatory estradiol has been reported to have a positive impact on conceptus development. Cows that exhibit estrus have been shown to have a longer conceptus length compared to no estrus cows on day 19 of gestation (Davoodi et al., 2016).

USE OF ESTRADIOL TO SYNCHRONIZE ESTRUS AND OVULATION

The ability of estradiol to synchronize standing estrus and ovulation has been the focus of many investigations. Using estradiol following PGF_{2α} administration increased expression of estrus (Cerri et al., 2004; Souza et al., 2005; Brusveen et al., 2009), reduced the interval to onset of estrus (Nancarrow and Radford, 1975; Ryan et al., 1995; Evans et al., 2003), induced a surge of LH (Short et al., 1973; Short et al., 1979), and reduced the interval from PGF_{2α} to ovulation (Ryan et al., 1995; Evans et al., 2003). However, the effect of supplemental estradiol on fertility during a synchronization protocol is not clear. When estradiol was administered at follicular wave emergence, fertility was reduced, but when administered when a dominant follicle was present on the ovary, it had no influence on fertility (Lane et al., 2001). Estradiol administered during follicular wave emergence reduced the maximum diameter of the ovulatory follicle, but estradiol administered when a dominant follicle was present did not affect maximum ovulatory follicle size (Evans et al., 2003). When estradiol was administered pre-breeding in sheep bred by natural mating or artificial insemination, there was no effect on pregnancy success (Inskeep et al., 1979). However, in the preceding studies all animals were inseminated following detection in standing estrus, and when cows were detected in standing estrus no differences were detected in circulating concentrations of estradiol (Perry et al., 2005). Administering estradiol before artificial insemination following a fixed-time insemination protocol improved fertility (Ahmadzadeh et al., 2003) when small follicles were induced to ovulate (Jinks et al., 2013). Time of estradiol administration and type of estradiol used can also influence fertility. Administering 1 mg of Estradiol Cypionate (ECP) did not induce an LH surge until 42 ± 2 hours and ovulation until 61 ± 3 hours after administration (Stevenson et al., 2004). However, administering 0.5 mg of estradiol benzoate induced an LH surge and ovulation 30 ± 3.1 and 72 ± 0.0 hours, respectively, after administration (Evans et al., 2003). Administration of estradiol-17β 8 hours (Souza et al., 2007) or 16 hours (Brusveen et al., 2009) before insemination did not improve pregnancy success and even tended to have a negative effect on fertility, but when

concentrations of estradiol were elevated for 1.9 days before insemination (follicle size held constant) pregnancy success was increased (Bridges et al., 2010). Therefore, the addition of estradiol to a synchronization protocol must consider dose, type of estrogen, and timing of administration in maximizing the benefits.

CONCLUSION

These studies indicate not only an importance in increasing preovulatory concentrations of estradiol but also the timing of the increase in estradiol and that increasing circulating concentrations of estradiol through supplementation may not be sufficient to increase fertility. Increased production of estradiol by the ovulatory follicle may be required to enhance fertility through the regulation of sperm transport, fertilization, oviductal secretions, uterine environment, and embryo survival.

Literature Cited

- Abe, H., Y. Sendai, T. Satoh, and H. Hoshi. 1995. Bovine oviduct-specific glycoprotein: a potent factor for maintenance of viability and motility of bovine spermatozoa in vitro. *Mol. Reprod. Dev.* 42: 226-232.
- Adashi, E. Y., and A. J. Hsueh. 1982. Estrogens augment the stimulation of ovarian aromatase activity by follicle-stimulating hormone in cultured rat granulosa cells. *J. Biol. Chem.* 257: 6077-6083.
- Ahmadzadeh, A., D. G. Falk, R. Manzo, C. B. Sellars, and J. C. Dalton. 2003. Effect of incorporation of a low dose of estradiol cypionate (ECP) into a timed artificial insemination protocol on estrous behavior and conception rates in beef cattle. *J. Anim. Sci.* 81(Suppl. 1): 180 (abstr M187).
- Albrecht, U. 2006. Orchestration of gene expression and physiology by the circadian clock. *J. Physiol Paris* 100: 243-251.
- Allison, A. J., and T. J. Robinson. 1972. The recovery of spermatozoa from the reproductive tract of the spayed ewe treated with progesterone and oestrogen. *J. Reprod. Fertil.* 31: 215-224.
- Allrich, R. D. 1994. Endocrine and neural control of estrus in dairy cows. *J. Dairy Sci.* 77: 2738-2744.
- Atkins, J. A., M. F. Smith, M. D. MacNeil, E. M. Jinks, F. M. Abreu, L. J. Alexander, and T. W. Geary. 2013. Pregnancy establishment and maintenance in cattle. *J. Anim. Sci.* 91: 722-733.
- Bartol, F. F., W. W. Thatcher, G. S. Lewis, E. L. Bliss, M. Drost, and F. W. Bazer. 1981. Effect of estradiol-17beta on PGF and total protein content in bovine uterine flushings and peripheral plasma concentration of 13, 14-dihydro-15-keto-PGF(2alpha). *Theriogenology* 15: 345-358.

- Bauersachs, S., S. E. Ulbrich, K. Gross, S. E. Schmidt, H. H. Meyer, R. Einspanier, H. Wenigerkind, M. Vermehren, H. Blum, F. Sinowatz, and E. Wolf. 2005. Gene expression profiling of bovine endometrium during the oestrous cycle: detection of molecular pathways involved in functional changes. *J. Mol. Endocrinol.* 34: 889-908.
- Bolzenius, J. K., R. A. Cushman, and G. A. Perry. 2016. Expression of Na⁺/H⁺ exchanger isoforms 1, 2, and 3 decreased after CIDR removal among cows that exhibited estrus, but did not change among cows that did not exhibit estrus, and as uterine pH decreased pregnancy success tended to increase. *Anim. Reprod. Sci.*: In Press.
- Bridges, G. A., M. L. Mussard, C. R. Burke, and M. L. Day. 2010. Influence of the length of proestrus on fertility and endocrine function in female cattle. *Anim. Reprod. Sci.* 117: 208-215.
- Bridges, G. A., M. L. Mussard, J. L. Pate, T. L. Ott, T. R. Hansen, and M. L. Day. 2012. Impact of preovulatory estradiol concentrations on conceptus development and uterine gene expression. *Anim. Reprod. Sci.* 133: 16-26.
- Britt, J. H. 1987. Induction and synchronization. In: E. S. E. Hafez (ed.) *Reproduction in Farm Animals*. Lea and Febiger, Philadelphia, PA.
- Brusveen, D. J., A. H. Souza, and M. C. Wiltbank. 2009. Effects of additional prostaglandin F2 α and estradiol-17 β during Ovsynch in lactating dairy cows. *J. Dairy Sci.* 92: 1412-1422.
- Buhi, W. C. 2002. Characterization and biological roles of oviduct-specific, oestrogen-dependent glycoprotein. *Reproduction* 123: 355-362.
- Burghardt, R. C., and E. Anderson. 1981. Hormonal modulation of gap junctions in rat ovarian follicles. *Cell Tissue Res.* 214: 181-193.
- Butler, W. R., J. J. Calaman, and S. W. Beam. 1996. Plasma and milk urea nitrogen in relation to pregnancy rate in lactating dairy cattle. *J. Anim. Sci.* 74: 858-865.
- Cerri, R. L., J. E. Santos, S. O. Juchem, K. N. Galvao, and R. C. Chebel. 2004. Timed artificial insemination with estradiol cypionate or insemination at estrus in high-producing dairy cows. *J. Dairy Sci.* 87: 3704-3715.
- Chabbert-Buffeta, N., D. C. Skinner, A. Caraty, and P. Bouchard. 2000. Neuroendocrine effects of progesterone. *Steroids* 65: 613-620.
- Chappell, P. E., R. S. White, and P. L. Mellon. 2003. Circadian gene expression regulates pulsatile gonadotropin-releasing hormone (GnRH) secretory patterns in the hypothalamic GnRH-secreting GT1-7 cell line. *J. Neurosci.* 23: 11202-11213.
- Chenault, J. R., W. W. Thatcher, P. S. Kalra, R. M. Abrams, and C. J. Wilcox. 1975. Transitory changes in plasma progestins, estradiol, and luteinizing hormone approaching ovulation in the bovine. *J. Dairy Sci.* 58: 709-717.
- Contri, A., A. Gloria, D. Robbe, C. Valorz, L. Wegher, and A. Carluccio. 2013. Kinematic study on the effect of pH on bull sperm function. *Anim. Reprod. Sci.* 136: 252-259.
- Dalton, J. C., S. Nadir, M. Noftsinger, J. Bame, and R. G. Saacke. 2006. Management factors associated with male fertility. In: *Applier Reproductive Strategies in Beef Cattle.*, Rapid City, SD. p 309-321.
- Davoodi, S., R. F. Cooke, A. C. Fernandes, B. I. Cappelozza, J. L. Vasconcelos, and R. L. Cerri. 2016. Expression of estrus modifies the gene expression profile in reproductive tissues on Day 19 of gestation in beef cows. *Theriogenology* 85: 645-655.

- DeJarnette, J. M., R. G. Saacke, J. Bame, and C. J. Vogler. 1992. Accessory sperm: their importance to fertility and embryo quality, and attempts to alter their numbers in artificially inseminated cattle. *J. Anim. Sci.* 70: 484-491.
- Diskin, M. G., and G. D. Niswender. 1989. Effect of progesterone supplementation on pregnancy and embryo survival in ewes. *J. Anim. Sci.* 67: 1559-1563.
- Dransfield, M. B., R. L. Nebel, R. E. Pearson, and L. D. Warnick. 1998. Timing of insemination for dairy cows identified in estrus by a radiotelemetric estrus detection system. *J. Dairy Sci.* 81: 1874-1882.
- Driancourt, M. A., B. Thuel, P. Mermillod, and P. Lonergan. 1998. Relationship between oocytes quality (measured after ivm, ivf and ivc of individual oocytes) and follicle function in cattle. *Therio* 49: 345 (abstract).
- Dupont, S., A. Krust, A. Gansmuller, A. Dierich, P. Chambon, and M. Mark. 2000. Effect of single and compound knockouts of estrogen receptors alpha (ERalpha) and beta (ERbeta) on mouse reproductive phenotypes. *Development* 127: 4277-4291.
- Elrod, C. C., and W. R. Butler. 1993. Reduction of fertility and alteration of uterine pH in heifers fed excess ruminally degradable protein. *J. Anim. Sci.* 71: 694-701.
- Elrod, C. C., M. Van Amburgh, and W. R. Butler. 1993. Alterations of pH in response to increased dietary protein in cattle are unique to the uterus. *J. Anim. Sci.* 71: 702-706.
- Erb, R. E., H. A. Garverick, R. D. Randel, B. L. Brown, and C. J. Callahan. 1976. Profiles of reproductive hormones associated with fertile and nonfertile inseminations of dairy cows. *Theriogenology* 5: 227-242.
- Evans, A. C., P. O'Keeffe, M. Mihm, J. F. Roche, K. L. Macmillan, and M. P. Boland. 2003. Effect of oestradiol benzoate given after prostaglandin at two stages of follicle wave development on oestrus synchronisation, the LH surge and ovulation in heifers. *Anim. Reprod. Sci.* 76: 13-23.
- Fanjul, L. F., C. M. Ruiz de Galarreta, and A. J. Hsueh. 1984. Estrogen regulation of progesterin biosynthetic enzymes in cultured rat granulosa cells. *Biol. Reprod.* 30: 903-912.
- Farookhi, R., and J. Desjardins. 1986. Luteinizing hormone receptor induction in dispersed granulosa cells requires estrogen. *Mol. Cell Endocrinol.* 47: 13-24.
- Fields, S. D., K. L. Gebhart, B. L. Perry, M. G. Gonda, C. L. Wright, R. C. Bott, and G. A. Perry. 2012. Influence of standing estrus before an injection of GnRH during a beef cattle fixed-time AI protocol on LH release, subsequent concentrations of progesterone, and steroidogenic enzyme expression. *Domest. Anim. Endocrinol.* 42: 11-19.
- Forde, N., M. E. Beltman, G. B. Duffy, P. Duffy, J. P. Mehta, P. O'Gaora, J. F. Roche, P. Lonergan, and M. A. Crowe. 2011. Changes in the endometrial transcriptome during the bovine estrous cycle: effect of low circulating progesterone and consequences for conceptus elongation. *Biol. Reprod.* 84: 266-278.
- Forde, N., F. Carter, T. Fair, M. A. Crowe, A. C. Evans, T. E. Spencer, F. W. Bazer, R. McBride, M. P. Boland, P. O'Gaora, P. Lonergan, and J. F. Roche. 2009. Progesterone-regulated changes in endometrial gene expression contribute to advanced conceptus development in cattle. *Biol. Reprod.* 81: 784-794.
- Fortune, J. E. 1986. Bovine theca and granulosa cells interact to promote androgen production. *Biol. Reprod.* 35: 292-299.
- Fortune, J. E., and S. M. Quirk. 1988. Regulation of steroidogenesis in bovine preovulatory follicles. *J. Anim. Sci.* 66: 1-8.

- Fortune, J. E., G. M. Rivera, A. C. Evans, and A. M. Turzillo. 2001. Differentiation of dominant versus subordinate follicles in cattle. *Biol. Reprod.* 65: 648-654.
- Fraser, L. R., G. Umar, and S. Sayed. 1993. Na(+)-requiring mechanisms modulate capacitation and acrosomal exocytosis in mouse spermatozoa. *J. Reprod. Fertil.* 97: 539-549.
- Geisert, R. D., G. L. Morgan, E. C. Short, Jr., and M. T. Zavy. 1992. Endocrine events associated with endometrial function and conceptus development in cattle. *Reprod. Fertil. Dev.* 4: 301-305.
- Ginther, O. J., M. C. Wiltbank, P. M. Fricke, J. R. Gibbons, and K. Kot. 1996. Selection of the dominant follicle in cattle. *Biol. Reprod.* 55: 1187-1194.
- Giroux-Widemann, V., P. Jouannet, I. Pignot-Paintrand, and D. Feneux. 1991. Effects of pH on the reactivation of human spermatozoa demembrated with Triton X-100. *Mol. Reprod. Dev.* 29: 157-162.
- Goldenberg, R. L., W. E. Bridson, and P. O. Kohler. 1972. Estrogen stimulation of progesterone synthesis by porcine granulosa cells in culture. *Biochem. Biophys. Res. Commun.* 48: 101-107.
- Goltz, J. S., T. K. Gardner, K. S. Kanous, and C. B. Lindemann. 1988. The interaction of pH and cyclic adenosine 3',5'-monophosphate on activation of motility in Triton X-100 extracted bull sperm. *Biol. Reprod.* 39: 1129-1136.
- Gray, C. A., K. M. Taylor, W. S. Ramsey, J. R. Hill, F. W. Bazer, F. F. Bartol, and T. E. Spencer. 2001. Endometrial glands are required for preimplantation conceptus elongation and survival. *Biol. Reprod.* 64: 1608-1613.
- Gregg, D. W., R. H. Schwall, and T. M. Nett. 1991. Regulation of gonadotropin secretion and number of gonadotropin-releasing hormone receptors by inhibin, activin-A, and estradiol. *Biol. Reprod.* 44: 725-732.
- Hasler, J. F. 2001. Factors affecting frozen and fresh embryo transfer pregnancy rates in cattle. *Theriogenology* 56: 1401-1415.
- Hawk, H. W. 1983. Sperm survival and transport in the female reproductive tract. *J. Dairy Sci.* 66: 2645-2660.
- Hawk, H. W., and T. Y. Tanabe. 1986. Effect of unilateral cornual insemination upon fertilization rate in superovulating and single-ovulating cattle. *J. Anim. Sci.* 63: 551-560.
- Hazeleger, N. L., D. J. Hill, R. B. Stubbings, and J. S. Walton. 1995. Relationship of morphology and follicular fluid environment of bovine oocytes to their development potential in vitro. *Therio* 43: 509-522.
- He, P. J., M. Hirata, N. Yamauchi, and M. A. Hattori. 2007. Up-regulation of Per1 expression by estradiol and progesterone in the rat uterus. *J. Endocrinol.* 194: 511-519.
- Henricks, D. M., D. R. Lamond, J. R. Hill, and J. F. Dickey. 1971. Plasma progesterone concentrations before mating and in early pregnancy in the beef heifer. *J. Anim. Sci.* 33: 450-454.
- Hill, J. L., S. K. Walker, G. H. Brown, and C. D. Nancarrow. 1996. The effects of an estrus-associated oviductal glycoprotein on the in vitro fertilization and development of ovine oocytes matured in vitro. *Theriogenology* 46: 1379-1388.
- Hunter, R. H., and I. Wilmut. 1984. Sperm transport in the cow: peri-ovulatory redistribution of viable cells within the oviduct. *Reprod. Nutr. Dev.* 24: 597-608.
- Ing, N. H., and R. M. Roberts. 1989. The major progesterone-modulated proteins secreted into the sheep uterus are members of the serpin superfamily of serine protease inhibitors. *J. Biol. Chem.* 264: 3372-3379.

- Inskeep, E. K., L. P. Stevens, and C. R. Rudy. 1979. Fertility in ewes receiving low doses of estradiol during synchronized estrus. *J. Anim. Sci.* 48: 52-53.
- Jinks, E. M., M. F. Smith, J. A. Atkins, K. G. Pohler, G. A. Perry, M. D. Macneil, A. J. Roberts, R. C. Waterman, L. J. Alexander, and T. W. Geary. 2013. Preovulatory estradiol and the establishment and maintenance of pregnancy in suckled beef cows. *J. Anim. Sci.*
- Johnson, K. R., R. H. Ross, and D. L. Fourn. 1958. Effect of progesterone administration on reproductive efficiency. *Theriogenology* 17: 386-390.
- Jones, R. L., C. Stoikos, J. K. Findlay, and L. A. Salamonsen. 2006. TGF-beta superfamily expression and actions in the endometrium and placenta. *Reproduction* 132: 217-232.
- Karsch, F. J., J. M. Bowen, A. Caraty, N. P. Evans, and S. M. Moenter. 1997. Gonadotropin-releasing hormone requirements for ovulation. *Biol. Reprod.* 56: 303-309.
- Kennaway, D. J. 2005. The role of circadian rhythmicity in reproduction. *Hum. Reprod. Update* 11: 91-101.
- Kerbler, T. L., M. M. Buhr, L. T. Jordan, K. E. Leslie, and J. S. Walton. 1997. Relationship between maternal plasma progesterone concentration and interferon-tau synthesis by the conceptus in cattle. *Therio* 47: 703-714.
- Kessel, B., Y. X. Liu, X. C. Jia, and A. J. Hsueh. 1985. Autocrine role of estrogens in the augmentation of luteinizing hormone receptor formation in cultured rat granulosa cells. *Biol. Reprod.* 32: 1038-1050.
- King, R. S., and G. J. Killian. 1994. Purification of bovine estrus-associated protein and localization of binding on sperm. *Biol. Reprod.* 51: 34-42.
- Lamming, G. E., and G. E. Mann. 1995. Control of endometrial oxytocin receptors and prostaglandin F_{2α} production in cows by progesterone and oestradiol. *J. Reprod. Fertil.* 103: 69-73.
- Lane, E. A., E. J. Austin, J. F. Roche, and M. A. Crowe. 2001. The effect of estradiol benzoate on synchrony of estrus and fertility in cattle after removal of a progesterone-releasing intravaginal device. *Theriogenology* 55: 1807-1818.
- Larimore, E. L., O. L. Amundson, S. L. Bird, B. J. Funnell, S. G. Kruse, G. A. Bridges, and G. A. Perry. 2015. Influence of estrus at fixed-time artificial insemination on early embryonic development in beef cattle. *J. Anim. Sci.* 93: 2806-2812.
- Looper, M. L., J. A. Vizcarra, R. P. Wettemann, J. R. Malayer, T. D. Braden, R. D. Geisert, and G. L. Morgan. 2003. Influence of estradiol, progesterone, and nutrition on concentrations of gonadotropins and GnRH receptors, and abundance of mRNA for GnRH receptors and gonadotropin subunits in pituitary glands of beef cows. *J. Anim. Sci.* 81: 269-278.
- Lopes, A. S., S. T. Butler, R. O. Gilbert, and W. R. Butler. 2007. Relationship of pre-ovulatory follicle size, estradiol concentrations and season to pregnancy outcome in dairy cows. *Anim. Reprod. Sci.* 99: 34-43.
- Lopez, H., L. D. Satter, and M. C. Wiltbank. 2004. Relationship between level of milk production and estrous behavior of lactating dairy cows. *Anim. Reprod. Sci.* 81: 209-223.
- Macmillan, K. L., and A. J. Peterson. 1993. A new intravaginal progesterone releasing device for cattle (CIDR-B) for oestrous synchronisation, increasing pregnancy rates and the treatment of post-partum anestrus. *Anim. Reprod. Sci.* 33: 1-25.
- Madsen, C. A., G. A. Perry, C. L. Mogck, R. F. Daly, M. D. MacNeil, and T. W. Geary. 2015. Effects of preovulatory estradiol on embryo survival and pregnancy establishment in beef cows. *Anim. Reprod. Sci.* 158: 96-103.

- Mann, G. E., and G. E. Lamming. 2001. Relationship between maternal endocrine environment, early embryo development and inhibition of the luteolytic mechanism in cows. *Reproduction* 121: 175-180.
- Mann, G. E., G. E. Lamming, and P. A. Fisher. 1998. Progesterone control of embryonic interferon tau production during early pregnancy in the cow. *J. Reprod. Fertil. Abstract series* 21: abstract 37.
- Mann, G. E., G. E. Lamming, R. S. Robinson, and D. C. Wathes. 1999. The regulation of interferon-tau production and uterine hormone receptors during early pregnancy. *J. Reprod. Fertil. Suppl.* 54: 317-328.
- McDonald, L. E., R. E. Nichols, and S. H. McNutt. 1952. Study of corpus luteum ablation and progesterone replacement therapy in cattle. *Am. J. Vet. Res.* 13: 446-451.
- McNatty, K. P., W. M. Hunter, A. S. MacNeilly, and R. S. Sawers. 1975. Changes in the concentration of pituitary and steroid hormones in the follicular fluid of human graafian follicles throughout the menstrual cycle. *J. Endocrinol.* 64: 555-571.
- McNatty, K. P., D. M. Smith, A. Makris, R. Osathanondh, and K. J. Ryan. 1979. The microenvironment of the human antral follicle: interrelationships among the steroid levels in antral fluid, the population of granulosa cells, and the status of the oocyte in vivo and in vitro. *J. Clin. Endocrinol. Metab.* 49: 851-860.
- Merk, F. B., C. R. Botticelli, and J. T. Albright. 1972. An intercellular response to estrogen by granulosa cells in the rat ovary; an electron microscope study. *Endocrinology* 90: 992-1007.
- Mermillod, P., B. Oussaid, and Y. Cognie. 1999. Aspects of follicular and oocyte maturation that affect the developmental potential of embryos. *J. Reprod. Fertil. Suppl.* 54: 449-460.
- Miller, B. G., and N. W. Moore. 1976. Effect of progesterone and oestradiol on endometrial metabolism and embryo survival in the ovariectomized ewe. *Theriogenology* 6: 636.
- Miller, B. G., N. W. Moore, L. Murphy, and G. M. Stone. 1977. Early pregnancy in the ewe: effects of oestradiol and progesterone on uterine metabolism and on embryo survival. *Aust. J. Biol. Sci.* 30: 279-288.
- Nadir, S., R. G. Saacke, J. Bame, J. Mullins, and S. Degelos. 1993. Effect of freezing semen and dosage of sperm on number of accessory sperm, fertility, and embryo quality in artificially inseminated cattle. *J. Anim. Sci.* 71: 199-204.
- Nakamura, T. J., T. Moriya, S. Inoue, T. Shimazoe, S. Watanabe, S. Ebihara, and K. Shinohara. 2005. Estrogen differentially regulates expression of Per1 and Per2 genes between central and peripheral clocks and between reproductive and nonreproductive tissues in female rats. *J Neurosci Res* 82: 622-630.
- Nancarrow, C. D., and H. M. Radford. 1975. Use of oestradiol benzoate to improve synchronization of oestrus in cattle. *J. Reprod. Fertil.* 43: 404.
- Nebel, R. L., and M. L. McGilliard. 1993. Interactions of high milk yield and reproductive performance in dairy cows. *J. Dairy Sci.* 76: 3257-3268.
- Nephew, K. P., H. Cardenas, K. E. McClure, T. L. Ott, F. W. Bazer, and W. F. Pope. 1994. Effects of administration of human chorionic gonadotropin or progesterone before maternal recognition of pregnancy on blastocyst development and pregnancy in sheep. *J. Anim. Sci.* 72: 453-458.
- Nett, T. M., A. M. Turzillo, M. Baratta, and L. A. Rispoli. 2002. Pituitary effects of steroid hormones on secretion of follicle-stimulating hormone and luteinizing hormone. *Domest. Anim. Endocrinol.* 23: 33-42.

- Noyes, R. W., C. E. Adams, and A. Walton. 1959. The passage of spermatozoa through the genital tract of female rabbits after ovariectomy and oestrogen treatment. *J. Endocrinol.* 18: 165-174.
- Ozturk, S., and R. Demir. 2010. Particular functions of estrogen and progesterone in establishment of uterine receptivity and embryo implantation. *Histol Histopathol* 25: 1215-1228.
- Parrish, J. J., J. L. Susko-Parrish, and N. L. First. 1989. Capacitation of bovine sperm by heparin: inhibitory effect of glucose and role of intracellular pH. *Biol. Reprod.* 41: 683-699.
- Parrott, J. A., and M. K. Skinner. 1998. Developmental and hormonal regulation of keratinocyte growth factor expression and action in the ovarian follicle. *Endocrinology* 139: 228-235.
- Peralta, O. A., R. E. Pearson, and R. L. Nebel. 2005. Comparison of three estrus detection systems during summer in a large commercial dairy herd. *Anim. Reprod. Sci.* 87: 59-72.
- Perry, G. A., and B. L. Perry. 2008a. Effect of preovulatory concentrations of estradiol and initiation of standing estrus on uterine pH in beef cows. *Domest. Anim. Endocrinol.* 34: 333-338.
- Perry, G. A., and B. L. Perry. 2008b. Effects of standing estrus and supplemental estradiol on changes in uterine pH during a fixed-time AI protocol. *J. Anim. Sci.* 86: 2928-2935.
- Perry, G. A., M. F. Smith, M. C. Lucy, J. A. Green, T. E. Parks, M. D. Macneil, A. J. Roberts, and T. W. Geary. 2005. Relationship between follicle size at insemination and pregnancy success. *Proc. Natl. Acad. Sci. U.S. A.* 102: 5268-5273.
- Perry, G. A., M. F. Smith, A. J. Roberts, M. D. MacNeil, and T. W. Geary. 2007. Relationship between size of ovulatory follicle and pregnancy success in beef heifers. *J. Anim. Sci.* 85: 684-689.
- Reilly, C. M., W. E. Cannady, V. B. Mahesh, V. S. Stopper, L. M. De Sevilla, and T. M. Mills. 1996. Duration of estrogen exposure prior to follicle-stimulating hormone stimulation is critical to granulosa cell growth and differentiation in rats. *Biol. Reprod.* 54: 1336-1342.
- Richardson, B. N., S. L. Hill, J. S. Stevenson, G. D. Djira, and G. A. Perry. 2016. Expression of Estrus before Fixed-time AI affects Conception Rates and Factors that Impact Expression of Estrus and the Repeatability of Expression of Estrus in Sequential Breeding Seasons. *Anim. Reprod. Sci.* <http://dx.doi.org/10.1016/j.anireprosci.2016.01.013>.
- Robinson, N. A., K. E. Leslie, and J. S. Walton. 1989. Effect of treatment with progesterone on pregnancy rate and plasma concentrations of progesterone in Holstein cows. *J. Dairy Sci.* 72: 202-207.
- Robinson, R. S., G. E. Mann, G. E. Lamming, and D. C. Wathes. 2001. Expression of oxytocin, oestrogen and progesterone receptors in uterine biopsy samples throughout the oestrous cycle and early pregnancy in cows. *Reproduction* 122: 965-979.
- Ryan, D. P., S. Snijders, A. Aarts, and K. J. O'Farrell. 1995. Effect of estradiol subsequent to induced luteolysis on development of the ovulatory follicle and interval to estrus and ovulation. *Theriogenology* 43: 310.
- Ryan, M., M. Mihm, and J. F. Roche. 1998. Effect of GnRH given before or after dominance on gonadotrophin response and fate of that follicle wave in postpartum dairy cows. *J. Reprod. Fertil.* 21: 61.
- Santos, J. E., W. W. Thatcher, R. C. Chebel, R. L. Cerri, and K. N. Galvao. 2004. The effect of embryonic death rates in cattle on the efficacy of estrus synchronization programs. *Anim. Reprod. Sci.* 82-83: 513-535.

- Seidel, G. E. 1995. Reproductive biotechnologies for profitable beef production. In: Proc. Beef Improvement Federation. : P 28 Sheridan, WY.
- Senger, P. L. 1994. The estrus detection problem: new concepts, technologies, and possibilities. *J. Dairy Sci.* 77: 2745-2753.
- Short, R. E., B. E. Howland, R. D. Randel, D. S. Christensen, and R. A. Bellows. 1973. Induced LH release in spayed cows. *J. Anim. Sci.* 37: 551-557.
- Short, R. E., R. D. Randel, R. B. Staigmiller, and R. A. Bellows. 1979. Factors affecting estrogen-induced LH release in the cow. *Biol. Reprod.* 21: 683-689.
- Souza, A. H., A. Gumen, E. P. Silva, A. P. Cunha, J. N. Guenther, C. M. Peto, D. Z. Caraviello, and M. C. Wiltbank. 2007. Supplementation with estradiol-17beta before the last gonadotropin-releasing hormone injection of the Ovsynch protocol in lactating dairy cows. *J. Dairy Sci.* 90: 4623-4634.
- Souza, A. H., A. Gumen, E. P. B. Silva, A. P. Cunha, J. N. Guenther, C. M. Peto, D. Z. Caraviello, and M. C. Wiltbank. 2005. Effect of estradiol-17B supplementation before the last GnRH of teh Ovsynch protocol in high producing dairy cows. *J. Anim. Sci.* 83(suppl. 1): 170 (abstr T157).
- Spencer, T. E., and F. W. Bazer. 2004. Uterine and placental factors regulating conceptus growth in domestic animals. *J. Anim. Sci.* 82 E-Suppl: E4-13.
- Spencer, T. E., O. Sandra, and E. Wolf. 2008. Genes involved in conceptus-endometrial interactions in ruminants: insights from reductionism and thoughts on holistic approaches. *Reproduction* 135: 165-179.
- Stauss, C. R., T. J. Votta, and S. S. Suarez. 1995. Sperm motility hyperactivation facilitates penetration of the hamster zona pellucida. *Biol. Reprod.* 53: 1280-1285.
- Stevenson, J. S., S. M. Tiffany, and M. C. Lucy. 2004. Use of estradiol cypionate as a substitute for GnRH in protocols for synchronizing ovulation in dairy cattle. *J. Dairy Sci.* 87: 3298-3305.
- Suarez, S. S., and X. Dai. 1992. Hyperactivation enhances mouse sperm capacity for penetrating viscoelastic media. *Biol. Reprod.* 46: 686-691.
- Sutton, R., C. D. Nancarrow, and A. L. C. Wallace. 1986. Oestrogen and seasonal effects on the production of an oestrus-associated glycoprotein in oviducal fluid of sheep. *J. Reprod. Fertil.* 77: 645-653.
- Teissier, M. P., H. Chable, S. Paulhac, and Y. Aubard. 2000. Comparison of follicle steroidogenesis from normal and polycystic ovaries in women undergoing IVF: relationship between steroid concentrations, follicle size, oocyte quality and fecundability. *Hum. Reprod.* 15: 2471-2477.
- Turzillo, A. M., C. E. Campion, C. M. Clay, and T. M. Nett. 1994. Regulation of gonadotropin-releasing hormone (GnRH) receptor messenger ribonucleic acid and GnRH receptors during the early preovulatory period in the ewe. *Endocrinology* 135: 1353-1358.
- Twagiramungu, H., L. A. Guilbault, and J. J. Dufour. 1995. Synchronization of ovarian follicular waves with a gonadotropin- releasing hormone agonist to increase the precision of estrus in cattle: a review. *J. Anim. Sci.* 73: 3141-3151.
- Ulbrich, S. E., T. Frohlich, K. Schulke, E. Englberger, N. Waldschmitt, G. J. Arnold, H. D. Reichenbach, M. Reichenbach, E. Wolf, H. H. Meyer, and S. Bauersachs. 2009. Evidence for estrogen-dependent uterine serpin (SERPINA14) expression during estrus in the bovine endometrial glandular epithelium and lumen. *Biol. Reprod.* 81: 795-805.

- Van Cleef, J., M. Drost, and W. W. Thatcher. 1991. Effects of postinsemination progesterone supplementation on fertility and subsequent estrous response on dairy heifers. *Theriogenology* 36: 795-807.
- Vasconcelos, J. L., R. Sartori, H. N. Oliveira, J. G. Guenther, and M. C. Wiltbank. 2001. Reduction in size of the ovulatory follicle reduces subsequent luteal size and pregnancy rate. *Theriogenology* 56: 307-314.
- Walton, J. S., G. W. Halbert, N. A. Robinson, and K. E. Leslie. 1990. Effects of progesterone and human chorionic gonadotrophin administration five days postinsemination on plasma and milk concentrations of progesterone and pregnancy rates of normal and repeat breeder dairy cows. *Can. J. Vet. Res.* 54: 305-308.
- Wang, X. F., M. K. Yu, S. Y. Lam, K. M. Leung, J. L. Jiang, P. S. Leung, W. H. Ko, P. Y. Leung, S. B. Chew, C. Q. Liu, C. M. Tse, and H. C. Chan. 2003. Expression, immunolocalization, and functional activity of Na⁺/H⁺ exchanger isoforms in mouse endometrial epithelium. *Biol. Reprod.* 68: 302-308.
- Wang, X. N., and G. S. Greenwald. 1993. Synergistic effects of steroids with FSH on folliculogenesis, steroidogenesis and FSH- and hCG-receptors in hypophysectomized mice. *J. Reprod. Fertil.* 99: 403-413.
- Welsh, T. H., Jr., L. Z. Zhuang, and A. J. Hsueh. 1983. Estrogen augmentation of gonadotropin-stimulated progesterin biosynthesis in cultured rat granulosa cells. *Endocrinology* 112: 1916-1924.
- Wise, M. E., D. Nieman, J. Stewart, and T. M. Nett. 1984. Effect of number of receptors for gonadotropin-releasing hormone on the release of luteinizing hormone. *Biol. Reprod.* 31: 1007-1013.
- Wong, P. Y., W. M. Lee, and A. Y. Tsang. 1981. The effects of extracellular sodium on acid release and motility initiation in rat caudal epididymal spermatozoa in vitro. *Exp. Cell Res.* 131: 97-104.
- Wu, J. C., S. C. Sealson, and W. L. Miller. 1994. Gonadal hormones and gonadotropin-releasing hormone (GnRH) alter messenger ribonucleic acid levels for GnRH receptors in sheep. *Endocrinology* 134: 1846-1850.
- Yanagimachi, R., and A. Bhattacharyya. 1988. Acrosome-reacted guinea pig spermatozoa become fusion competent in the presence of extracellular potassium ions. *J. Exp. Zool.* 248: 354-360.
- Zelinski, M. B., N. A. Hirota, E. J. Keenan, and F. Stormshak. 1980. Influence of exogenous estradiol-17 beta on endometrial progesterone and estrogen receptors during the luteal phase of the ovine estrous cycle. *Biol. Reprod.* 23: 743-751.
- Zeng, Y., J. A. Oberdorf, and H. M. Florman. 1996. pH regulation in mouse sperm: identification of Na⁽⁺⁾-, Cl⁽⁻⁾-, and HCO₃⁽⁻⁾-dependent and arylaminobenzoate-dependent regulatory mechanisms and characterization of their roles in sperm capacitation. *Dev. Biol.* 173: 510-520.
- Zhuang, L. Z., E. Y. Adashi, and A. J. Hsueh. 1982. Direct enhancement of gonadotropin-stimulated ovarian estrogen biosynthesis by estrogen and clomiphene citrate. *Endocrinology* 110: 2219-2221.